

COMBINATORIAL K -THEORY

ANDERS SKOVSTED BUCH

1. INTRODUCTION

Let X be a non-singular quasi-projective variety. The *Grothendieck ring* $K(X)$ of X is defined as the free abelian group generated by isomorphism classes $[F]$ of algebraic vector bundles on X , modulo the relations $[F] = [F'] + [F'']$ whenever there exists a short exact sequence $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$. Multiplication is given by the tensor product, $[F_1] \cdot [F_2] = [F_1 \otimes F_2]$.

If $Y \subset X$ is a closed subvariety, we can find a locally free resolution

$$0 \rightarrow F_n \rightarrow F_{n-1} \rightarrow \cdots \rightarrow F_1 \rightarrow F_0 \rightarrow \mathcal{O}_Y \rightarrow 0$$


of the structure sheaf of Y . The *Grothendieck class* of Y is defined as

$$[\mathcal{O}_Y] = \sum_{i=0}^n (-1)^i [F_i] \in K(X).$$

Suppose Z is another closed subvariety of X . In intersection theory one studies the geometry of the intersection $Y \cap Z$. This can be done using (Chow) cohomology by examining the product $[Y] \cdot [Z] \in H^*(X)$, or with K -theory by studying the product of Grothendieck classes $[\mathcal{O}_Y] \cdot [\mathcal{O}_Z] \in K(X)$. When Y and Z meet sufficiently transversally, these products give the respective classes of the intersection, in the sense that $[Y \cap Z] = [Y] \cdot [Z] \in H^*(X)$ and $[\mathcal{O}_{Y \cap Z}] = [\mathcal{O}_Y] \cdot [\mathcal{O}_Z] \in K(X)$.

The Grothendieck ring has a *topological filtration* $K(X) = \mathcal{F}_0 \supset \mathcal{F}_1 \supset \cdots$ by ideals. The ideal \mathcal{F}_i is generated by all classes $[\mathcal{O}_Y]$ for which $Y \subset X$ has codimension at least i . There is a group homomorphism $H^*(X) \rightarrow \text{gr}(K(X)) = \bigoplus_{i \geq 0} \mathcal{F}_i / \mathcal{F}_{i+1}$ from the Chow ring to the associated graded ring of $K(X)$ given by $[Y] \mapsto [\mathcal{O}_Y]$, and this map becomes an isomorphism of rings after tensoring with \mathbb{Q} (see [10, Ex. 15.2.16]). The geometric meaning of this is that intersection theory in the cohomology ring only captures the lowest graded piece of information available in the Grothendieck ring.

Example 1. Let $C \subset \mathbb{P}^2$ be a curve of degree 3. Since C can be degenerated into a union of three lines, its cohomology class is $[C] = 3[\text{line}]$. The Grothendieck class is finer and can “see” that these lines overlap in three points. Thus $[\mathcal{O}_C] = 3[\mathcal{O}_{\text{line}}] - 3[\mathcal{O}_{\text{point}}]$. This example generalizes easily to a hypersurface of any degree in \mathbb{P}^n .



Date: April 2003.

The goal of these notes is to explain the combinatorics that arises in the study of K -theoretic intersection theory of some concrete varieties and degeneracy loci. The material is mostly taken from the papers [4, 3], as well as from joint work [7] with A. Kresch, H. Tamvakis, and A. Yong.

Our notes are based on a lecture series presented at the *Mini-School on Schubert Varieties* at the Banach Center in the Spring of 2003. We thank Piotr Pragacz and Andrzej Weber, and the Banach Center for their kind invitation and hospitality. We also thank R. Vakil for private communication concerning a symmetry property of K -theoretic Schubert structure constants.

2. K -THEORY OF GRASSMANNIANS

Let $X = \text{Gr}(d, n)$ be the Grassmann variety of d -dimensional subspaces of \mathbb{C}^n . For each partition $\lambda = (\lambda_1 \geq \dots \geq \lambda_d \geq 0)$ with $\lambda_1 \leq n - d$, there is a Schubert variety

$$X_\lambda = \{V \in X \mid \dim(V \cap \mathbb{C}^{n-d+i-\lambda_i}) \geq i \forall 1 \leq i \leq d\}.$$

Here $\mathbb{C}^k \subset \mathbb{C}^n$ denotes the subspace of vectors where the last $n - k$ coordinates are zero. The codimension of X_λ in X is equal to the weight $|\lambda| = \sum_{i=1}^d \lambda_i$ of λ . We will identify the partition λ with its *Young diagram* of boxes, which has λ_1 boxes in the top row, λ_2 boxes in the second row, etc. If we let $R = (n - d)^d$ denote a rectangular partition with d rows and $n - d$ columns, then the Schubert varieties in X correspond to the Young diagrams λ contained in R .

$$R = \begin{array}{c} \boxed{\lambda} \\ \text{---} \\ \boxed{} \end{array} \begin{array}{c} d \\ \text{---} \\ n - d \end{array}$$

We will write $\mathcal{O}_\lambda = [\mathcal{O}_{X_\lambda}]$ for the Grothendieck class of the Schubert variety X_λ . These classes form a basis for the Grothendieck ring of X :

$$K(X) = \bigoplus_{\lambda \subset R} \mathbb{Z} \cdot \mathcal{O}_\lambda.$$

It is interesting to note that, as abstract rings, the Grothendieck and cohomology rings of a Grassmann variety are isomorphic. Despite this fact, the K -theoretic intersection theory has a richer structure than the cohomological intersection theory.

Given two partitions $\lambda, \mu \subset R$, we can write

$$\mathcal{O}_\lambda \cdot \mathcal{O}_\mu = \sum_{\nu} c_{\lambda\mu}^\nu \mathcal{O}_\nu$$

for unique structure constants $c_{\lambda\mu}^\nu \in \mathbb{Z}$. Our first goal will be to describe these constants, which determine the K -theoretic intersection theory on X . Using the isomorphism $H^*(X) \cong \text{gr}(K(X))$ it follows that the integer $c_{\lambda\mu}^\nu$ is non-zero only when $|\nu| \geq |\lambda| + |\mu|$, and when we have equality $|\nu| = |\lambda| + |\mu|$, the coefficient $c_{\lambda\mu}^\nu$ is a structure constant in the cohomology ring. These cohomology constants are called *Littlewood-Richardson coefficients* and play an important role in numerous fields, including representation theory, geometry, and combinatorics, see e.g. [12]. Lenart [21] has proved that the K -theoretic Pieri coefficients $c_{(p),\mu}^\nu$ can be expressed as plus or minus a binomial coefficient.

Set-valued tableaux. We will formulate a general rule for the structure constants $c_{\lambda\mu}^\nu$, and for this purpose we introduce some notation. Define a *set-valued tableau* to be a labeling of the boxes of a skew diagram with finite non-empty sets of positive integers, so that the numbers in the diagram increase weakly along rows and increase strictly down columns. More precisely, the maximum number of any box must be less than or equal to the minimum of the box to the right of it, and strictly smaller than the minimum of the box below it. For example

$$T = \begin{array}{ccc} & & \boxed{1} \ \boxed{23} \\ & \boxed{12} \ \boxed{234} & \\ \boxed{2} \ \boxed{35} \ \boxed{7} & & \end{array}$$

is a set-valued tableau whose shape is the skew diagram $(4, 3, 3)/(2, 1)$. Define the *content* of a set-valued tableau to be the sequence (c_1, c_2, \dots) , where c_i is the number of boxes containing the integer i . We also define the *word* of a set-valued tableau to be the sequence of integers in its boxes when read from left to right then bottom to top; the integers of a single box are arranged in increasing order. The above tableau T has content $(2, 4, 3, 1, 1, 0, 1)$ and word $(2, 3, 5, 7, 1, 2, 2, 3, 4, 1, 2, 3)$. A sequence of integers is called a *reverse lattice word* if every integer $i \geq 2$ in the sequence is followed by more $i - 1$'s than i 's.

Given two partitions λ and μ , we let $\lambda * \mu$ denote the skew shape obtained by attaching the Young diagrams for λ and μ corner to corner as shown.

$$\lambda * \mu = \begin{array}{c} \mu \\ \lambda \end{array}$$

The following theorem from [4] is a K -theoretic generalization of the classical Littlewood-Richardson rule [22].

Theorem 1. *The structure constant $c_{\lambda\mu}^\nu$ is equal to $(-1)^{|\nu|-|\lambda|+|\mu|}$ times the number of set-valued tableaux of shape $\lambda * \mu$, such that the word is a reverse lattice word with content ν .*


Example 2. Let $\lambda = \mu = (2, 1)$ and $\nu = (3, 3, 1)$. Then there are two set-valued tableaux of shape $\lambda * \mu$ such that the word is a reverse lattice word with content ν .

$$\begin{array}{ccc} & & \boxed{1} \ \boxed{1} \\ & & \boxed{2} \\ \boxed{1} \ \boxed{23} & & \\ \boxed{2} & & \end{array} \quad \begin{array}{ccc} & & \boxed{1} \ \boxed{1} \\ & & \boxed{2} \\ \boxed{1} \ \boxed{2} & & \\ \boxed{23} & & \end{array}$$

It follows that $c_{\lambda\mu}^\nu = -2$.

It follows from the theorem that K -theoretic structure constants on Grassmannians have signs that alternate with codimension, that is $(-1)^{|\nu|-|\lambda|+|\mu|} c_{\lambda\mu}^\nu \geq 0$. This alternation in sign has in fact been established by Brion for all homogeneous spaces G/P , where G is a simply connected semisimple algebraic group [2]. One can also derive from Theorem 1 that, if \mathcal{O}_ν occurs with non-zero coefficient in $\mathcal{O}_\lambda \cdot \mathcal{O}_\mu$, then we have $X_\lambda \cap X_\mu \supset X_\nu \supset \bigcap_\rho X_\rho$, where the intersection is over all partitions ρ such that $[X_\rho]$ occurs in the cohomology product $[X_\lambda] \cdot [X_\mu] \in H^*(X)$.

Another consequence of the theorem is the following symmetry property of the K -theoretic Schubert structure constants $c_{\lambda\mu}^\nu$. For a partition $\lambda \subset R$ we let $\lambda^\vee = (n - d - \lambda_d, \dots, n - d - \lambda_1)$ be the dual partition.



Corollary 1. *For any three partitions $\lambda, \mu, \nu \subset R$ we have $c_{\lambda\mu}^\nu = c_{\lambda^\vee\mu^\vee}^{\nu^\vee}$.*

Notice that when $|\nu| = |\lambda| + |\mu|$, this is immediate from the identity $c_{\lambda\mu}^\nu = \int_X [X_\lambda] \cdot [X_\mu] \cdot [X_{\nu^\vee}]$ for cohomological structure constants.

The proof given below was produced in response to a question of R. Vakil, who had observed that the corollary follows from a conjectured K -theoretic puzzle rule of A. Knutson and T. Tao. We note that this conjecture has now been proved by Knutson and Vakil by translating K -theoretic puzzles to set-valued tableaux [16].

Proof. Let $\rho : X \rightarrow \{\text{point}\}$ be a map to a point. The corresponding pushforward map $\rho_* : K(X) \rightarrow K(\text{point}) = \mathbb{Z}$ is given by $\rho_*(\mathcal{O}_\lambda) = 1$ for all $\lambda \subset R$. Define the bilinear form

$$K(X) \otimes K(X) \rightarrow \mathbb{Z} \quad ; \quad \alpha \otimes \beta \mapsto \rho_*(\alpha \cdot \beta).$$

We claim that this is a perfect pairing, and that the dual basis element of \mathcal{O}_λ equals $t\mathcal{O}_{\lambda^\vee}$, where $t = 1 - \mathcal{O}_{(1)}$. (This class t represents the top exterior power of the tautological subbundle on X .) The corollary is immediate from this claim since $c_{\lambda\mu}^\nu = \rho_*(\mathcal{O}_\lambda \cdot \mathcal{O}_\mu \cdot t\mathcal{O}_{\nu^\vee}) = c_{\lambda^\vee\mu^\vee}^{\nu^\vee}$.

It follows from Theorem 1 that $t\mathcal{O}_\lambda = \mathcal{O}_\lambda - \mathcal{O}_{(1)} \cdot \mathcal{O}_\lambda = \sum_\nu (-1)^{|\nu/\lambda|} \mathcal{O}_\nu$, where the sum is over all partitions ν such that $\lambda \subset \nu \subset R$ and ν/λ has at most one box in any row or column. This implies that $\rho_*(t\mathcal{O}_\lambda) = \delta_{\lambda,R}$, and we obtain

$$\rho_*(t\mathcal{O}_\lambda \cdot \mathcal{O}_\mu) = \sum_\nu c_{\lambda\mu}^\nu \rho_*(t\mathcal{O}_\nu) = c_{\lambda\mu}^R.$$

Finally, it is an easy exercise to check from the theorem that $c_{\lambda\mu}^R = \delta_{\lambda,\mu^\vee}$. \square

3. THE BIALGEBRA OF STABLE GROTHENDIECK POLYNOMIALS

For a set-valued tableau T with content (c_1, c_2, \dots, c_r) , we set $x^T = x_1^{c_1} x_2^{c_2} \dots x_r^{c_r}$. For any partition λ we define a *stable Grothendieck polynomial* for this partition by

$$G_\lambda = \sum_{\text{sh}(T)=\lambda} (-1)^{|T| - |\lambda|} x^T \in \mathbb{Z}[[x_1, x_2, \dots]].$$

The sum is over all set-valued tableau T of shape λ . This is a special case of the stable Grothendieck polynomials studied by Fomin and Kirillov [8] (see section 6). One can check from the definition that the power series G_λ is symmetric in the variables x_i . Furthermore, the term of lowest degree in G_λ is the Schur function for λ . In particular, the elements G_λ are linearly independent. We let Γ denote their linear span.

$$\Gamma = \bigoplus_{\lambda} \mathbb{Z} G_\lambda \subset \mathbb{Z}[[x_1, x_2, \dots]]$$

If $E = L_1 \oplus \cdots \oplus L_e$ is a vector bundle on a variety X , which can be written as a direct sum of line bundles L_i , we define

$$G_\lambda(E) = G_\lambda(1 - L_1^{-1}, \dots, 1 - L_e^{-1}, 0, 0, \dots) \in K(X).$$

Since G_λ is symmetric, this can be written as a polynomial in the exterior powers of the dual bundle E^\vee , so the definition also makes sense when E is not a direct sum of line bundles. When $X = \text{Gr}(d, \mathbb{C}^n)$ is a Grassmann variety and $S \subset \mathbb{C}^n \times X$ the tautological subbundle on X , it follows from [20] or [13, Thm. 3] that $\mathcal{O}_\lambda = G_\lambda(S^\vee)$. Theorem 1 generalizes to the following statement [4, Cor. 5.5].

Theorem 2. *For any partitions λ and μ we have*

$$G_\lambda \cdot G_\mu = \sum c_{\lambda\mu}^\nu G_\nu$$

where the sum is over all partitions ν , and the constants $c_{\lambda\mu}^\nu$ are given by Theorem 1.

It follows from this theorem that each product $G_\lambda \cdot G_\mu$ is a finite linear combination of stable Grothendieck polynomials G_ν . In particular, Γ is a subring of the power series ring $\mathbb{Z}[[x_1, x_2, \dots]]$.

For partitions λ , μ , and ν , we set $d_{\lambda\mu}^\nu = c_{\nu R}^\rho$, where R is a rectangular partition which is wider than λ and taller than μ , and $\rho = (R + \mu, \lambda)$ is the partition obtained by attaching λ and μ to the sides of R .

$$\rho = (R + \mu, \lambda) = \begin{array}{c} \boxed{R} \quad \mu \\ \lambda \end{array}$$

It is not hard to see that $d_{\lambda\mu}^\nu$ is independent of the choice of the rectangle R . Given a finite set of variables x_1, \dots, x_n , we write $G_\lambda(x_1, \dots, x_n) = G_\lambda(x_1, \dots, x_n, 0, 0, \dots)$. The coefficients $d_{\lambda\mu}^\nu$ have the following interpretation.

Theorem 3. *For any partition ν and integers $0 < m < n$ we have*

$$G_\nu(x_1, \dots, x_m, x_{m+1}, \dots, x_n) = \sum_{\lambda, \mu} d_{\lambda\mu}^\nu G_\lambda(x_1, \dots, x_m) \cdot G_\mu(x_{m+1}, \dots, x_n).$$

Define a coproduct $\Delta : \Gamma \rightarrow \Gamma \otimes \Gamma$ by

$$\Delta(G_\nu) = \sum_{\lambda, \mu} d_{\lambda\mu}^\nu G_\lambda \otimes G_\mu.$$

This coproduct corresponds to the pullback along the embedding

$$\text{Gr}(d_1, n_1) \times \text{Gr}(d_2, n_2) \rightarrow \text{Gr}(d_1 + d_2, n_1 + n_2)$$

which maps a pair $(V_1 \subset \mathbb{C}^{n_1}, V_2 \subset \mathbb{C}^{n_2})$ to $V_1 \oplus V_2 \subset \mathbb{C}^{n_1+n_2}$.

It follows from theorems 2 and 3 that the coproduct Δ makes Γ into a commutative and cocommutative bialgebra. The linear map $\Gamma \rightarrow \mathbb{Z}$ which sends $1 = G_{(0)}$ to one and G_λ to zero for $|\lambda| > 0$ furthermore gives a counit. This bialgebra is a K -theoretic parallel of the ring of symmetric functions.

The bialgebra Γ has an involution $\Gamma \rightarrow \Gamma$ defined by $G_\lambda \mapsto G_{\lambda'}$, where λ' is the conjugate partition of λ . This involution corresponds to the duality isomorphism $\text{Gr}(d, n) \cong \text{Gr}(n - d, n)$ of Grassmannians. Equivalently, the structure constants of Γ satisfy the identities $c_{\lambda\mu}^\nu = c_{\lambda'\mu'}^{\nu'}$ and $d_{\lambda\mu}^\nu = d_{\lambda'\mu'}^{\nu'}$.

4. GEOMETRIC SPECIALIZATIONS OF STABLE GROTHENDIECK POLYNOMIALS

Given two vector bundles E and F over the variety X and a partition ν , we define

$$G_\nu(F - E) = \sum_{\lambda, \mu} d_{\lambda\mu}^\nu G_\lambda(F) \cdot G_{\mu'}(E^\vee) \in K(X)$$

where E^\vee is the dual bundle of E . It is proved in [4] that $G_\lambda(F - E)$ is a specialization of the double stable Grothendieck polynomials of Fomin and Kirillov [8]. It therefore follows from a super symmetry property of the latter polynomials that, for any vector bundle H we have

$$(1) \quad G_\lambda(F \oplus H - E \oplus H) = G_\lambda(F - E).$$

In other words, G_λ gives a well defined linear map $G_\lambda : K(X) \rightarrow K(X)$. Since the coproduct Δ is coassociative, we furthermore deduce that

$$(2) \quad G_\nu(F - E) = \sum_{\lambda, \mu} d_{\lambda\mu}^\nu G_\lambda(F - H) \cdot G_\mu(H - E).$$

Another useful identity is $G_\lambda(F - E) = G_{\lambda'}(E^\vee - F^\vee)$, which was first proved by Fomin (see [4, Lemma 3.4].)

If the bundles E and F have ranks e and f , and if λ and μ are partitions such that $\ell(\lambda) \leq f$ and $\mu_1 \leq e$, we also have the factorization formula

$$(3) \quad G_{(e^f)+\lambda, \mu}(F - E) = G_\lambda(F) G_{(e^f)}(F - E) G_\mu(-E)$$

where $(e^f) + \lambda, \mu$ denotes the partition $(e + \lambda_1, \dots, e + \lambda_f, \mu_1, \mu_2, \dots)$.

More generally, the following Gysin formula was proved in [3]. It generalizes Pragacz's cohomological Gysin formula from [24] and [14, (4.4)] to K -theory.

Theorem 4. *Let E and F be vector bundles of ranks e and f over X , and let $\pi : \text{Gr}(d, F) \rightarrow X$ denote the Grassmann bundle of subbundles of F of rank d , with universal exact sequence $0 \rightarrow S \rightarrow \pi^* F \rightarrow Q \rightarrow 0$. Set $q = f - d$. Let λ and μ be partitions such that $\ell(\lambda) \leq q$, and $\lambda_q \geq \max(e, \mu_1 + d)$. Then we have in $K(X)$ that*

$$\pi_*(G_\lambda(Q - \pi^* E) \cdot G_\mu(S - \pi^* E)) = G_{(\lambda_1 - d, \dots, \lambda_q - d, \mu_1, \mu_2, \dots)}(F - E).$$

Like Pragacz's cohomological Gysin formula, this result remains true with slightly weaker conditions. Namely, one may drop the requirement that $\lambda_q - d \geq \mu_1$. This means that the sequence $I = (\lambda_1 - d, \dots, \lambda_q - d, \mu_1, \mu_2, \dots)$ is not necessarily a partition; the definition of G_I when I is an arbitrary sequence of integers is given [3, §3], but will not be discussed here. It is curious to note that the same formula is also true when λ and μ are arbitrary partitions such that $\ell(\lambda) \leq q$ and $\ell(\mu) \leq d$ [24], [14, (4.3)], but this case of the formula is false in K -theory.

5. DEGENERACY LOCI

Let $\phi : E \rightarrow F$ be a map of vector bundles of ranks e and f over X . Given an integer $t \leq \min(e, f)$, there is a degeneracy locus

$$\Omega_t = \Omega_t(\phi) = \{x \in X \mid \text{rank}(\phi_x : E(x) \rightarrow F(x)) \leq t\}.$$

This locus has a natural structure of subscheme of X , given as the zero-section of the map $\wedge^{t+1} \phi : \wedge^{t+1} E \rightarrow \wedge^{t+1} F$. The expected (and maximal possible) codimension of Ω_t is the number $(e - t)(f - t)$. When this codimension is attained, the

classical Thom-Porteous formula expresses the cohomology class of Ω_t as a Schur determinant. This formula generalizes to K -theory as follows.

Theorem 5. *If Ω_t has codimension $(e-t)(f-t)$ in X then its Grothendieck class is given by*

$$[\mathcal{O}_{\Omega_t}] = G_{(e-t)f-t}(F - E) \in K(X).$$

The Thom-Porteous degeneracy locus Ω_t can be generalized as follows. Let $E_0 \rightarrow E_1 \rightarrow \cdots \rightarrow E_n$ be a sequence of vector bundles and bundle maps over X . A set of *rank conditions* for this sequence is a collection $r = \{r_{ij}\}$ of non-negative integers, for $0 \leq i < j \leq n$. This data defines the *quiver variety*

$$\Omega_r = \Omega_r(E_\bullet) = \{x \in X \mid \text{rank}(E_i(x) \rightarrow E_j(x)) \leq r_{ij} \forall i < j\}.$$

We will demand that the rank conditions can occur, i.e. there exists a sequence of vector spaces and linear maps $V_0 \rightarrow V_1 \rightarrow \cdots \rightarrow V_n$ such that $\dim(V_i) = \text{rank}(E_i)$ and $\text{rank}(V_i \rightarrow V_j) = r_{ij}$ for all $i < j$. In other words, the rank conditions should correspond to an orbit of a quiver representation as in [1]. In this case, the expected codimension of the quiver variety Ω_r is equal to

$$d(r) = \sum_{i < j} (r_{i,j-1} - r_{ij})(r_{i+1,j} - r_{ij})$$

where we set $r_{ii} = \text{rank}(E_i)$ for all i . In work with Fulton [6], we proved a formula for the cohomology class of a quiver variety Ω_r . The following theorem from [3] generalizes this formula to K -theory.

Theorem 6. *Let $E_0 \rightarrow E_1 \rightarrow \cdots \rightarrow E_n$ be a sequence of vector bundle maps such that the quiver variety $\Omega_r(E_\bullet)$ has codimension $d(r)$ in X . Then the Grothendieck class of this quiver variety is given by*

$$[\mathcal{O}_{\Omega_r}] = \sum c_\mu(r) G_{\mu_1}(E_1 - E_0) \cdot G_{\mu_2}(E_2 - E_1) \cdots G_{\mu_n}(E_n - E_{n-1}) \in K(X)$$

where the sum is over sequences $\mu = (\mu_1, \dots, \mu_n)$ of partitions μ_i , and the coefficients $c_\mu(r)$ are integers given by the rank conditions.

The coefficients $c_\mu(r)$ of this formula are called *quiver coefficients*. They are uniquely determined by the fact that the theorem is true for all varieties X , as well as the requirement that $c_\mu(r)$ is non-zero only if $\ell(\mu_i) \leq r_{ii}$ for all i . These coefficients can furthermore be computed by an explicit combinatorial algorithm based on the operations of the bialgebra Γ , see [3, §4].

A quiver coefficient $c_\mu(r)$ is non-zero only when $\sum |\mu_i| \geq d(r)$. When $\sum |\mu_i| = d(r)$, the coefficient $c_\mu(r)$ also appears in the formula for the cohomology class of Ω_r ; these coefficients are called *cohomological quiver coefficients*. It was conjectured in [6] and proved by Knutson, Miller, and Shimozono in [15] that cohomological quiver coefficients are non-negative. The K -theoretic quiver coefficients were conjectured to have alternating signs, in the following sense.¹

Conjecture 1. *For any rank conditions r and sequence of partitions μ , we have*

$$(-1)^{\sum |\mu_i| - d(r)} c_\mu(r) \geq 0.$$

¹After these lectures were presented, this conjecture from [3] has been proved in the papers [5] and [23].

We explain how to prove Theorem 6 for a sequence of three vector bundles $E \rightarrow F \rightarrow H$ of ranks e, f , and h . In this case the quiver variety is defined as

$$\Omega = \{x \in X \mid \text{rank}(E \rightarrow F) \leq r, \text{rank}(F \rightarrow H) \leq s, \text{rank}(E \rightarrow H) \leq t\}$$

for some non-negative integers r, s , and t . The requirement that these rank conditions can occur says that $r \leq \min(e, f)$, $s \leq \min(f, h)$, $t \leq \min(r, s)$, and $f + t \geq r + s$.

To make sure that the bundle maps are maximally generic, we start by replacing X with the bundle $\text{Hom}(E, F) \oplus \text{Hom}(F, H)$ and the bundle maps with the universal maps $E \rightarrow F \rightarrow H$ on this bundle. (In the rest of this section we avoid explicit notation for pullbacks of bundles.) Then form the product $Y = \text{Gr}(r, F) \times_X \text{Gr}(s, H)$ of Grassmann bundles with projection $\pi : Y \rightarrow X$, and let $F' \subset F$ and $H' \subset H$ denote the tautological subbundles on Y . On the subset $Z = Z(E \rightarrow F/F') \cap Z(F \rightarrow H/H') \subset Y$, the bundle map from E to F factors through F' , and the map $F \rightarrow H$ factors through H' . This gives a map $F' \rightarrow F \rightarrow H'$ on Z , and we let $\Omega_t = \Omega_t(F' \rightarrow H') \subset Z$ denote the corresponding Thom-Porteous locus.

Since all the rank conditions are satisfied on Ω_t , it follows that $\pi(\Omega_t) \subset \Omega$. In fact, by using that the bundle maps are universal, one can show that π maps Ω_t birationally onto Ω (see [6, Lemma 1].) Since quiver varieties have rational singularities [17], it follows that

$$\pi_*([\mathcal{O}_{\Omega_t}]) = [\mathcal{O}_{\Omega}] \in K(X).$$

Let $R = (r-t)^{s-t}$ be a rectangle with $s-t$ rows and $r-t$ columns. By Theorem 5 and equations (2) and (3) we have

$$\begin{aligned} [\mathcal{O}_{\Omega_t}] &= G_{(e)f-r}(F/F' - E) G_{(f)h-s}(H/H' - F) G_R(H' - F') \\ &= G_{(e)f-r}(F/F' - E) G_{(f)h-s}(H/H' - F) \sum_{\lambda, \mu} d_{\lambda, \mu}^R G_{\lambda}(F/F') G_{\mu}(H' - F) \\ &= \sum_{\lambda, \mu} d_{\lambda, \mu}^R G_{(e)f-r+\lambda}(F/F' - E) G_{(f)h-s}(H/H' - F) G_{\mu}(H' - F) \end{aligned}$$

in the Grothendieck ring of Y . Using Theorem 4 we therefore obtain

$$[\mathcal{O}_{\Omega}] = \pi_*([\mathcal{O}_{\Omega_t}]) = \sum_{\lambda, \mu} d_{\lambda, \mu}^R G_{(e-r)f-r+\lambda}(F - E) \cdot G_{(f-s)h-s, \mu}(H - F)$$

in $K(X)$ as required. Notice that the quiver coefficients for a sequence of three bundles are just the coproduct coefficients $d_{\lambda, \mu}^R$. In particular, they have alternating signs.

6. GROTHENDIECK POLYNOMIALS

For each permutation $w \in S_{n+1}$, there is a *double Grothendieck polynomial* $\mathfrak{G}_w(x; y) = \mathfrak{G}_w(x_1, \dots, x_n; y_1, \dots, y_n)$ defined as follows [20, 18]. If $w = w_0$ is the longest permutation in S_{n+1} , then we set

$$\mathfrak{G}_{w_0}(x; y) = \prod_{i+j \leq n+1} (x_i + y_j - x_i y_j).$$

Otherwise there is a simple transposition $s_i \in S_{n+1}$ such that $\ell(ws_i) = \ell(w) + 1$. In this case we set

$$\mathfrak{G}_w(x; y) = \pi_i(\mathfrak{G}_{ws_i})$$

where the *Demazure operator* π_i is defined by

$$\pi_i(f) = \frac{(1 - x_{i+1})f(x_1, \dots, x_i, x_{i+1}, \dots, x_n) - (1 - x_i)f(x_1, \dots, x_{i+1}, x_i, \dots, x_n)}{x_i - x_{i+1}}.$$

Grothendieck polynomials were introduced by Lascoux and Schützenberger [20, 18] as representatives for the Grothendieck classes of Schubert varieties in a flag variety. More generally, the double Grothendieck polynomials have the following interpretation. Let $F_1 \subset F_2 \subset \dots \subset F_n \rightarrow H_n \rightarrow \dots \rightarrow H_2 \rightarrow H_1$ be a sequence of bundles on X consisting of a full flag F_\bullet with a general map to a dual full flag H_\bullet , such that $\text{rank}(F_i) = \text{rank}(H_i) = i$ for all i . Given a permutation $w \in S_{n+1}$, Fulton [11] defined the degeneracy locus

$$\Omega_w = \Omega_w(F_\bullet \rightarrow H_\bullet) = \{x \in X \mid \text{rank}(F_q(x) \rightarrow H_p(x)) \leq r_w(p, q) \forall p, q\},$$

where $r_w(p, q) = \#\{i \leq p \mid w(i) \leq q\}$. This type of degeneracy locus generalizes Schubert varieties in flag varieties. In fact, if $X = \text{Fl}(n+1)$ is the variety of full flags in \mathbb{C}^{n+1} , and if $F_\bullet \subset \mathbb{C}^{n+1} \times X$ denotes the tautological flag on X and $\mathbb{C}^{n+1} \rightarrow H_\bullet$ is a fixed dual flag, then $\Omega_w \subset X$ is a Schubert variety.

Fulton proved that the cohomology class of Ω_w is given by a double Schubert polynomial [11]. This was generalized to the following formula for the Grothendieck class in [3, Thm. 2.1] as an application of [13, Thm. 3].

Theorem 7. *If the codimension of Ω_w equals the length of w , then*

$$[\mathcal{O}_{\Omega_w}] = \mathfrak{G}_w(x; y)$$

where $x_i = 1 - L_i^{-1} \in K(X)$ for $L_i = \ker(H_i \rightarrow H_{i-1})$, and $y_i = 1 - M_i \in K(X)$ for $M_i = F_i/F_{i-1}$.

Fulton's degeneracy locus Ω_w is a special case of a quiver variety, that is $\Omega_w = \Omega_r(F_\bullet \rightarrow H_\bullet)$ for a set of rank conditions r depending on the permutation w . It follows from this that $\mathfrak{G}_w(x; y) = [\mathcal{O}_{\Omega_r}]$, so

$$(4) \quad \mathfrak{G}_w(x; y) = \sum_{\mu} c_{w, \mu} G_{\mu_1}(F_2 - F_1) \cdots G_{\mu_n}(H_n - F_n) \cdots G_{\mu_{2n-1}}(H_1 - H_2)$$

where the constants $c_{w, \mu} = c_{\mu}(r)$ are quiver coefficients. Note that $G_{\mu_i}(F_{i+1} - F_i) = G_{\mu_i}(M_{i+1})$ is either zero or a power of $1 - M_{i+1}^{-1} = 1 - (1 - y_{i+1})^{-1}$, which can be expressed as a formal power series in y_{i+1} . Similarly $G_{\mu_{2n-i}}(H_i - H_{i+1}) = G_{\mu_{2n-i}}(-L_{i+1})$ is a power of $1 - (1 - x_{i+1})^{-1}$.

The double Grothendieck polynomials are related to stable Grothendieck polynomials as follows. Given a permutation $w \in S_{n+1}$, we let $1^m \times w \in S_{m+n+1}$ denote the shifted permutation, which acts as the identity on the set $\{1, 2, \dots, m\}$ and maps $m+i$ to $m+w(i)$ for $i \geq 1$. The *stable Grothendieck polynomial* $G_w(x; y)$ of Fomin and Kirillov [8] is defined by

$$G_w(x_1, \dots, x_m; y_1, \dots, y_m) = \mathfrak{G}_{1^m \times w}(x_1, \dots, x_m, 0, \dots, 0; y_1, \dots, y_m, 0, \dots, 0).$$

Let $\lambda = (\lambda_1 \geq \dots \geq \lambda_k \geq 0)$ be a partition. The corresponding Grassmannian permutation w_λ with descent at position k is defined by $w_\lambda(i) = i + \lambda_{k+1-i}$ for $i \leq k$ and $w_\lambda(i) < w_\lambda(i+1)$ for $i \neq k$. With the notation of section 3, we have $G_\lambda(x) = G_{w_\lambda}(x; 0)$ and

$$G_{w_\lambda}(x; y) = G_\lambda(x; y) = \sum_{\sigma, \tau} d_{\sigma\tau}^\lambda G_\sigma(x) G_{\tau'}(y).$$

Every stable Grothendieck polynomial $G_w(x; y)$ can be written as a finite linear combination of the polynomials $G_\lambda(x; y)$ given by partitions [4]. We will describe a formula of Lascoux for the coefficients of this linear combination, which shows that they have alternating signs.

Given a permutation w , let r be the last descent position of w , i.e. r is maximal such that $w(r) > w(r+1)$. Set $w' = w\tau_{rk}$ where $k > r$ is maximal such that $w(r) > w(k)$. We also set $I(w) = \{i < r \mid \ell(w'\tau_{ir}) = \ell(w)\}$. Define a relation \triangleright on the set of all permutations as follows. If $I(w) = \emptyset$ we write $w \triangleright v$ if and only if $v = 1 \times w$. Otherwise we write $w \triangleright v$ if and only if there exist elements $i_1 < \dots < i_p$ of $I(w)$, $p \geq 1$, such that $v = w'\tau_{i_1 r} \dots \tau_{i_p r}$. The following is an immediate consequence of [19, Thm. 4].

Theorem 8 (Lascoux). *For any permutation w we have*

$$G_w = \sum_{\lambda} a_{w,\lambda} G_{\lambda}$$

where the sum is over all partitions λ , and $a_{w,\lambda}$ is equal to $(-1)^{|\lambda| - \ell(w)}$ times the number of sequences $w = w_1 \triangleright w_2 \triangleright \dots \triangleright w_m$ such that $w_m = w_{\lambda}$ is a Grassmannian permutation for λ and w_i is not Grassmannian for $i < m$.

7. ALTERNATING SIGNS OF THE COEFFICIENTS $c_{w,\mu}$

In this section we outline a proof that the quiver coefficients $c_{w,\mu}$ of (4) have alternating signs, based on our joint paper [7] with Kresch, Tamvakis, and Yong.

Suppose we are given a sequence of vector bundles

$$F_1 \subset F_2 \subset \dots \subset F_n \subset V \rightarrow H_n \rightarrow \dots \rightarrow H_2 \rightarrow H_1$$

consisting of a full flag F_{\bullet} of a bundle V of rank $n+1$, followed by a dual full flag H_{\bullet} of V . Set $F'_i = V/F_i$ and $H'_i = \ker(V \rightarrow H_i)$. We then obtain a sequence

$$H'_n \subset \dots \subset H'_1 \subset V \rightarrow F'_1 \rightarrow \dots \rightarrow F'_n,$$

and it is an easy exercise to show that $\Omega_w(F_{\bullet} \rightarrow H_{\bullet}) = \Omega_{\hat{w}}(H'_{\bullet} \rightarrow F'_{\bullet})$ as subschemes of X , where $\hat{w} = w_0 w^{-1} w_0$. It follows that $\mathfrak{G}_w(x; y) = [\mathcal{O}_{\Omega_{\hat{w}}}]$, so we obtain

$$\begin{aligned} \mathfrak{G}_w(x; y) &= \sum_{\mu} c_{\hat{w},\mu} G_{\mu_1}(H'_{n-1} - H'_n) \cdots G_{\mu_n}(F'_1 - H'_1) \cdots G_{\mu_{2n-1}}(F'_n - F'_{n-1}) \\ &= \sum_{\mu} c_{\hat{w},\mu} G_{\mu_1}(x_n) \cdots G_{\mu_n}(x_1; y_1) \cdots G_{\mu_{2n-1}}(0; y_n) \end{aligned}$$

where x_i and y_i are defined as in Theorem 7. Notice that $G_{\lambda}(x_i)$ equals x_i^p when $\lambda = (p)$ is a single row with p boxes, and is zero otherwise. Similarly $G_{\lambda}(0; y_i)$ is a power of y_i or zero, and furthermore $G_{\lambda}(x_1; y_1) = \sum_{\sigma,\tau} d_{\sigma\tau}^{\lambda} G_{\sigma}(x_1) G_{\tau'}(y_1)$.

Corollary 2. *The monomial coefficients of Grothendieck polynomials are special cases of the quiver coefficients $c_{w,\mu}$.*

The *degenerate Hecke algebra* is the free \mathbb{Z} -algebra \mathcal{H} generated by symbols s_1, s_2, \dots , modulo the relations $s_i^2 = s_i$ and $s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}$ for all i , and $s_i s_j = s_j s_i$ for $|i - j| \geq 2$. This algebra has a basis of permutations, corresponding to reduced expressions in the generators.

Now define the *universal Grothendieck polynomial* for the permutation $w \in S_n$ to be the element

$$P_w = \sum_{\mu} c_{w,\mu} G_{\mu_1} \otimes \cdots \otimes G_{\mu_{2n-1}} \in \Gamma^{\otimes 2n-1}.$$

The following theorem gives an explicit formula for these polynomials.

Theorem 9. *For $w \in S_{n+1}$ we have*

$$P_w = \sum (-1)^{\sum \ell(u_i) - \ell(w)} G_{u_1} \otimes G_{u_2} \otimes \cdots \otimes G_{u_{2n-1}} \in \Gamma^{\otimes 2n-1},$$

where the sum is over all factorizations $w = u_1 \cdot u_2 \cdots u_{2n-1}$ in the degenerate Hecke algebra \mathcal{H} such that $u_i \in S_{\min(i, 2n-i)+1}$ for each i .

This theorem combined with Lascoux's formula for the expansion of stable Grothendieck polynomials in the basis of Γ implies the following explicit formula for the quiver coefficients $c_{w,\mu}$.

Corollary 3. *The quiver coefficients $c_{w,\mu}$ of P_w are given by*

$$c_{w,\mu} = (-1)^{\sum |\mu_i| - \ell(w)} \sum \prod_{i=1}^{2n-1} |a_{u_i, \mu_i}|,$$

where the sum is over all factorizations $w = u_1 \cdot u_2 \cdots u_{2n-1}$ in the degenerate Hecke algebra \mathcal{H} such that $u_i \in S_{\min(i, 2n-i)+1}$ for each i , and the constants a_{u_i, μ_i} are given by Theorem 8.

The proof of Theorem 9 is based on some identities of universal Grothendieck polynomials. For $1 \leq i \leq j \leq 2n-1$ we define

$$P_w[i, j] = \sum_{\mu: \mu_k = \emptyset \text{ for } k \notin [i, j]} c_{w,\mu} 1 \otimes \cdots \otimes 1 \otimes G_{\mu_i} \otimes \cdots \otimes G_{\mu_j} \otimes 1 \otimes \cdots \otimes 1.$$

Using a Cauchy identity for double Grothendieck polynomials of Fomin and Kirillov [9] as well as some geometry of quiver varieties, one obtains the identity

$$P_w = \sum_{u \cdot v = w \in \mathcal{H}} (-1)^{\ell(uv)} P_u[1, i] \cdot P_v[i+1, 2n-1].$$

By iterating this formula, we obtain

$$P_w = \sum_{u_1 \cdots u_{2n-1} = w \in \mathcal{H}} (-1)^{\sum \ell(u_i) - \ell(w)} P_{u_1}[1, 1] \cdot P_{u_2}[2, 2] \cdots P_{u_{2n-1}}[2n-1, 2n-1].$$

Finally we use the identity

$$P_w[i, i] = \begin{cases} 1^{\otimes i-1} \otimes G_w \otimes 1^{\otimes 2n-1-i} & \text{if } w \in S_{\min(i, 2n-i)+1} \\ 0 & \text{otherwise.} \end{cases}$$

This was proved in [3] for $i = n$, and the remaining cases are easy consequences of this. Theorem 9 follows immediately from these identities.

REFERENCES

- [1] S. Abeasis and A. Del Fra, *Degenerations for the representations of an equioriented quiver of type A_m* , Boll. Un. Mat. Ital. Suppl. (1980), no. 2, 157–171. MR **84e**:16019
- [2] M. Brion, *Positivity in the Grothendieck group of complex flag varieties*, J. Algebra **258** (2002), no. 1, 137–159, Special issue in celebration of Claudio Procesi's 60th birthday. MR **2003m**:14017
- [3] A. S. Buch, *Grothendieck classes of quiver varieties*, Duke Math. J. **115** (2002), no. 1, 75–103. MR **2003m**:14018

- [4] ———, *A Littlewood-Richardson rule for the K -theory of Grassmannians*, Acta Math. **189** (2002), no. 1, 37–78. MR **2003j**:14062
- [5] ———, *Alternating signs of quiver coefficients*, preprint, 2003.
- [6] A. S. Buch and W. Fulton, *Chern class formulas for quiver varieties*, Invent. Math. **135** (1999), no. 3, 665–687. MR **2000f**:14087
- [7] A. S. Buch, A. Kresch, H. Tamvakis, and A. Yong, *Grothendieck polynomials and quiver formulas*, To appear in Amer. J. Math., 2003.
- [8] S. Fomin and A. N. Kirillov, *Grothendieck polynomials and the Yang-Baxter equation*, Proc. Formal Power Series and Alg. Comb. (1994), 183–190.
- [9] ———, *The Yang-Baxter equation, symmetric functions, and Schubert polynomials*, Discrete Math. **153** (1996), 123–143.
- [10] W. Fulton, *Intersection theory*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 2, Springer-Verlag, Berlin, 1984. MR **85k**:14004
- [11] ———, *Flags, Schubert polynomials, degeneracy loci, and determinantal formulas*, Duke Math. J. **65** (1992), no. 3, 381–420. MR **93e**:14007
- [12] ———, *Young tableaux*, London Mathematical Society Student Texts, vol. 35, Cambridge University Press, Cambridge, 1997, With applications to representation theory and geometry. MR **99f**:05119
- [13] W. Fulton and A. Lascoux, *A Pieri formula in the Grothendieck ring of a flag bundle*, Duke Math. J. **76** (1994), no. 3, 711–729. MR **96j**:14036
- [14] W. Fulton and P. Pragacz, *Schubert varieties and degeneracy loci*, Lecture Notes in Mathematics, vol. 1689, Springer-Verlag, Berlin, 1998, Appendix J by the authors in collaboration with I. Ciocan-Fontanine. MR **99m**:14092
- [15] A. Knutson, E. Miller, and M. Shimozono, *Four positive formulas for type A quiver polynomials*, preprint, 2003.
- [16] A. Knutson and R. Vakil, manuscript in preparation.
- [17] V. Lakshmibai and P. Magyar, *Degeneracy schemes, quiver schemes, and Schubert varieties*, Internat. Math. Res. Notices (1998), no. 12, 627–640. MR **99g**:14065
- [18] A. Lascoux, *Anneau de Grothendieck de la variété de drapeaux*, The Grothendieck Festschrift, Vol. III, Progr. Math., vol. 88, Birkhäuser Boston, Boston, MA, 1990, pp. 1–34. MR **92j**:14064
- [19] ———, *Transition on Grothendieck polynomials*, Physics and combinatorics, 2000 (Nagoya), World Sci. Publishing, River Edge, NJ, 2001, pp. 164–179. MR **2002k**:14082
- [20] A. Lascoux and M.-P. Schützenberger, *Structure de Hopf de l’anneau de cohomologie et de l’anneau de Grothendieck d’une variété de drapeaux*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), no. 11, 629–633. MR **84b**:14030
- [21] C. Lenart, *Combinatorial aspects of the K -theory of Grassmannians*, Ann. Comb. **4** (2000), no. 1, 67–82. MR **2001j**:05124
- [22] D. E. Littlewood and A. R. Richardson, *Group characters and algebra*, Phil. Trans. R. Soc., A **233** (1934), 99–141.
- [23] E. Miller, *Alternating formulae for K -theoretic quiver polynomials*, preprint, 2003.
- [24] P. Pragacz, *Enumerative geometry of degeneracy loci*, Ann. Sci. École Norm. Sup. (4) **21** (1988), no. 3, 413–454. MR **90e**:14004