

Modulated elliptic wave and a train of asymptotic solitons in a vicinity of the leading edge for MKdV

Alexander Minakov

SISSA

via Bonomea 265

34136 Trieste

minakov.ilt@gmail.com

We discuss the behavior of the solution of the modified Korteweg-de Vries equation of a step type in a vicinity of the leading edge.

The initial data tends to zero as $x \rightarrow +\infty$ and to some positive constant c as $x \rightarrow -\infty$. In 1989 E. Khruslov and V. Kotlyarov have found that for a large time the solution breaks up into a train of asymptotic solitons located in the domain $4c^2t - C_N \log t < x < 4c^2t$ (C_N is a constant). The number N of these solitons grows unboundedly as $t \rightarrow +\infty$.

The behavior of the solution in the domain slightly apart from the leading edge $(-6c^2 + \varepsilon)t < x < (4c^2 - \varepsilon)t$ was studied in [3], and this solution is described by a modulated elliptic wave.

It would be natural to assume that the modulated elliptic asymptotics is still valid in the domain $(4c^2 - \varepsilon)t < x < 4c^2t$, but in [4] it was shown that the modulated elliptic wave and the train of asymptotic solitons do not coincide in the domain $4c^2t - C_N \log t < x < 4c^2t$ because of a phase shift. This indicates a possible presence of a further transition region in the already small domain $(4c^2 - \varepsilon)t < x < 4c^2t - C_N \log t$.

References

- [1] E. Ya. Khruslov: *Asymptotics of the solution of the Cauchy problem for the Korteweg-de Vries equation with initial data of step type*, Matem. Sbornik (New Series) 99(141):2 (1976), 261–281.
- [2] E. Ya. Khruslov, V. P. Kotlyarov: *Asymptotic solitons of the modified Korteweg-de Vries equation*, Inverse problems 5/6 (1989), 1075-1088.
- [3] V. Kotlyarov, A. Minakov: *Riemann-Hilbert problem to the modified Korteweg-de Vries equation: long-time dynamics of the steplike initial data*, J. Math. Phys 51, no.9 (2010), 093506, 31pp.
- [4] V. Kotlyarov, A. Minakov: *Modulated elliptic wave and asymptotic solitons in a shock problem to the modified Korteweg-de Vries equation*, J. Phys. A 13 (2015), 305201, 35pp.